

**ARYA GROUP OF COLLEGES**  
**I MID TERM EXAMINATION 2018-19 (I Sem.)**  
**1FY2-01\_Engineering Mathematics- I**  
**BRANCH: Common to All**

**Max Marks:- 80****Time:- 2 hrs.****PART A (Attempt All)**

- Q.1 (a) If  $u = \tan^{-1} \left( \frac{x^2+y^2}{x-y} \right)$ , prove that (i)  $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = \sin 2u$ . 5\*4
- (b) If  $\vec{q} = (y+z)\hat{i} + (z+x)\hat{j} + (x+y)\hat{k}$ , then show that  $\vec{q}$  is irrotational.
- (c) Define Beta & Gamma function
- (d) Find the equation of tangent plane & normal line to the surface  $x^2 + y^2 + z - 9 = 0$ , at P (1,2,4).
- (e) Find the stationary point of the following function:  $u(x, y) = x^2 + y^2 + \frac{2}{x} + \frac{2}{y}$

**PART B (Attempt any Four)**

- Q.2 (a) If  $u = f \left( \frac{x}{y}, \frac{y}{z}, \frac{z}{x} \right)$ ; then show that  $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} + z \frac{\partial u}{\partial z} = 0$
- (b) Prove that  $B(m, n) = \frac{\Gamma m \Gamma n}{\Gamma(m+n)}$  4\*8
- (c) Find the divergence & curl of the vector  $\vec{F} = \nabla(x^3 + y^3 + z^3 - 3xyz)$
- (d) If  $u = f(r)$ , where  $r^2 = x^2 + y^2$  prove that
- $$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = f''(r) + f'(r)/r$$
- (e) Find the extreme value of  $x^2 + y^2 + z^2$  subject to the condition  $ax + by + cz = p$ .
- (f) If  $f(x, y) = \begin{cases} \frac{x^2 - y^2}{x^2 + y^2}; & \text{when } x \neq 0, y \neq 0 \\ 0; & \text{when } x = y = 0 \end{cases}$  then discuss the continuity of  $f(x, y)$  at the origin.

**PART C (Attempt any Two)**

- Q.3 (a) Find the volume & surface area of solid generated by revolving the curve astroid  $x = a \cos^3 t, y = a \sin^3 t$  about x axis.
- (b) (i) Evaluate  $I = \int_0^\infty \frac{dx}{1+x^4}$       (ii)  $I = \int_0^{\pi/6} \cos^4 \theta \sin^2 6\theta d\theta$  2\*14
- (c) If  $u = f(x, y)$  where  $x = \psi \cos \alpha - n \sin \alpha, y = \psi \sin \alpha + n \cos \alpha$   
then prove that  $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = \frac{\partial^2 u}{\partial \psi^2} + \frac{\partial^2 u}{\partial n^2}$ ,  
 $\alpha$  is a constant.

$$2x =$$

$$2(x^2 + y^2 + z^2) + (ax + by + cz)\lambda = 0$$

$$2f + p\lambda = 0$$

$$\boxed{\lambda = -\frac{2f}{p}}$$

$$x = -\frac{ax}{2}$$

$$x = a \cos^3 t$$

$$-\frac{a}{2} x - \frac{2f}{p}$$

$$\frac{dx}{dt} = 3a \cos^2 t$$

$$-\frac{a(x^2 + y^2 + z^2)}{(ax + by + cz)}$$

$$-\underline{\sin^5}$$

$$2f + p\lambda = 0$$

$$\boxed{\lambda = -\frac{2f}{p}}$$

$$2 \times \int_0^{r_2} \pi y^2 \frac{dx}{dt} dt$$

$$x = -\frac{ax}{2} \quad y = -\frac{by}{2} \quad z = -\frac{cz}{2}$$

$$-\frac{a^2}{2}\lambda - \frac{b^2}{2}\lambda - \frac{c^2}{2}\lambda = p$$

$$\lambda = \frac{-2p}{a^2 + b^2 + c^2}$$

[Section-A] (Start to write From here)

Q. 11. @ If  $u = \tan^{-1} \left\{ \frac{x^3 + y^3}{x - y} \right\}$  Prove that (i)  $\frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} = \sin 2u$

Sol. Given  $u = \tan^{-1} \left\{ \frac{x^3 + y^3}{x - y} \right\}$  Then  $\tan u = f(u) = \frac{x^3 + y^3}{x - y} = \frac{x^3}{x - y} \left( 1 + \left( \frac{y}{x} \right)^3 \right)$

$f(u) = \tan u = x^2 \not\propto \left( \frac{y}{x} \right)$  Here  $f(u) = \tan u$  is the homogeneous

function of degree  $n=2$ , we have by Production of a Euler's Theorem.

(i)  $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = g(u) = n f(u) = 2 \tan u = 2 \sin u \cdot \cos^2 u$

$\frac{\partial u}{\partial x} = \frac{\cos u}{\sin u} = \cot u$  Ans.

(2)


$$\nabla \cdot (\vec{F} + \vec{G}) = (\vec{F} + \vec{G}) \cdot \nabla = (\vec{F} \cdot \nabla) + (\vec{G} \cdot \nabla)$$

SP40 RINGO INDUSTRIAL AREA DELHI ROAD RAIPUR  
MANUFACTURED BY TATA TRUCKS

**Sol:** we have **LITERMATIC** is non-singular vector field.

$$\text{curl } \vec{F} = 0$$

$$0 \times \vec{F} = 0$$



$$\text{curl } \vec{F} = \hat{i} \left( \frac{\partial}{\partial y} (x+y) - \frac{\partial}{\partial z} (x+z) \right) - \hat{j} \left( \frac{\partial}{\partial x} (x+y) - \frac{\partial}{\partial z} (y+z) \right) + \hat{k} \left( \frac{\partial}{\partial x} (x+z) - \frac{\partial}{\partial y} (y+z) \right)$$

$$\nabla \times \vec{F} = \hat{i} (1-1) - \hat{j} (1-1) + \hat{k} (1-1)$$

$$= \hat{o}_i - \hat{o}_j + \hat{o}_k = 0$$

Hence  $\vec{F}$  is non-singular vector field

Pranav

(2)

(C) Define Beta and Gamma function.

Beta function :-

if it is denoted by  $B(m, n)$  and define

$$\text{as } B(m, n) = \int_0^1 x^{m-1} (1-x)^{n-1} dx \quad m > 0 \quad n > 0$$

Gamma function :-

it is denoted by  $\Gamma(m)$

$$\text{and define as } \Gamma(m) = \int_0^\infty e^{-x} x^{m-1} dx \quad m > 0$$

all the functions are the type of Improper Integrals.

Ans.

(d)

Find the equations of Tangent plane and Normal line  
to the surface.

$$\phi = x^2 + y^2 + z - 9 = 0 \quad \text{at } P(1, 2, 4)$$

Sol:- Given  $\phi = x^2 + y^2 + z - 9 = 0$

we have eqn of Tangent plane.

$$(x-x_0) f_x(P) + (y-y_0) f_y(P) + (z-z_0) f_z(P) = 0 \quad \text{--- (1)}$$

and Normal line:-

$$\frac{x-x_0}{f_x(P)} = \frac{y-y_0}{f_y(P)} = \frac{z-z_0}{f_z(P)} = t \quad \text{--- (2)}$$

Param of w.r.t x, y and z.

$$\frac{\partial \phi}{\partial x} = 2x, \quad \frac{\partial \phi}{\partial y} = 2y, \quad \frac{\partial \phi}{\partial z} = 1$$

$$f_x(P) = 2, \quad f_y(P) = 4, \quad f_z(P) = 1$$

True Normal line:-

$$\left[ \begin{array}{l} x-1 \\ 2 \\ 2 \end{array} \right] = \left[ \begin{array}{l} y-2 \\ 4 \\ 1 \end{array} \right] = \left[ \begin{array}{l} z-4 \\ -1 \\ 1 \end{array} \right]$$

Tangent plane:-

$$(x-1)\cdot 2 + (y-2)\cdot 4 + (z-4)\cdot 1 = 0$$

$$\left[ \begin{array}{l} 2x+4y+z=14 \end{array} \right]$$

An.

Q(1)(e)

Find the stationary point of the following function

$$u(x,y) = x^2 + y^2 + \frac{2}{x} + \frac{2}{y} \quad \text{--- (1)}$$

Sol:- Given

$$u(x,y) = x^2 + y^2 + \frac{2}{x} + \frac{2}{y}$$

(i) Find  $\frac{\partial u}{\partial x}$  and  $\frac{\partial u}{\partial y}$

$$\frac{\partial u}{\partial x} = 2x - \frac{2}{x^2} \quad \text{and} \quad \frac{\partial u}{\partial y} = 2y - \frac{2}{y^2}$$

(ii) equating  $\frac{\partial u}{\partial x} = 0$  and  $\frac{\partial u}{\partial y} = 0$

$$\text{thus } 2x - 2 = 0 \quad \text{--- (1)} \quad \text{and} \quad 2y - \frac{2}{y^2} = 0 \quad \text{--- (2)}$$

$$x^3 - 1 = 0$$

$$(x-1) = 0$$

$$\text{thus} \quad x=1 \quad y=1$$

i.e. The stationary point is  $(1,1)$  at which we check, the

Maxima and Minima of the function

Ans.

[Part-B]

Q. 2 (a)

If  $u = f\left(\frac{x}{y}, \frac{y}{z}, \frac{z}{x}\right)$  Then show that-

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} + z \frac{\partial u}{\partial z} = 0$$

Sol:-

Given  $u = f\left(\frac{x}{y}, \frac{y}{z}, \frac{z}{x}\right)$  let  $t_1 = \frac{x}{y}$

$u = f(t_1, t_2, t_3)$  and  $t_1, t_2, t_3$  are the function of  $x, y$  and  $z$ .

$$t_3 = \frac{z}{x}$$

Now  $\frac{\partial t_1}{\partial x} = \frac{1}{y}$   $\frac{\partial t_1}{\partial y} = -\frac{x}{y^2}$   $\frac{\partial t_1}{\partial z} = 0$

$\frac{\partial t_2}{\partial x} = 0$   $\frac{\partial t_2}{\partial y} = \frac{1}{z}$   $\frac{\partial t_2}{\partial z} = -\frac{y}{z^2}$

$\frac{\partial t_3}{\partial x} = -\frac{z}{x^2}$   $\frac{\partial t_3}{\partial y} = 0$   $\frac{\partial t_3}{\partial z} = \frac{1}{x}$

We have by Total derivative.

$$\frac{\partial y}{\partial x} = \frac{\partial y}{\partial t_1} \frac{\partial t_1}{\partial x} + \frac{\partial y}{\partial t_2} \frac{\partial t_2}{\partial x} + \frac{\partial y}{\partial t_3} \frac{\partial t_3}{\partial x}$$

$$= \frac{\partial y}{\partial t_1} \left( \frac{1}{y} \right) + \frac{\partial y}{\partial t_2} \times 0 + \frac{\partial y}{\partial t_3} \times \left( -\frac{z}{x^2} \right) \quad \text{--- (1)} \times x$$

$$\frac{\partial y}{\partial x} = \frac{\partial y}{\partial t_1} \frac{\partial t_1}{\partial y} + \frac{\partial y}{\partial t_2} \frac{\partial t_2}{\partial y} + \frac{\partial y}{\partial t_3} \frac{\partial t_3}{\partial y}$$

$$= \frac{\partial y}{\partial t_1} \left( -\frac{x}{y} \right) + \frac{\partial y}{\partial t_2} \left( \frac{1}{z} \right) + \frac{\partial y}{\partial t_3} \times 0 \quad \text{--- (2)} \times y$$

$$\frac{\partial y}{\partial z} = \frac{\partial y}{\partial t_1} \frac{\partial t_1}{\partial z} + \frac{\partial y}{\partial t_2} \frac{\partial t_2}{\partial z} + \frac{\partial y}{\partial t_3} \frac{\partial t_3}{\partial z}$$

$$= \frac{\partial y}{\partial t_1} \times 0 + \frac{\partial y}{\partial t_2} \times -\frac{y}{z} + \frac{\partial y}{\partial t_3} \times \frac{1}{x} \quad \text{--- (3)} \times z.$$

Adding (1) (2) and (3)

$$x \frac{\partial y}{\partial x} + y \frac{\partial y}{\partial y} + z \frac{\partial y}{\partial z} = \frac{x}{y} \frac{\partial y}{\partial t_1} - \frac{z}{x} \frac{\partial y}{\partial t_3} + \left( -\frac{x}{y} \right) \frac{\partial y}{\partial t_1} + \frac{y}{z} \frac{\partial y}{\partial t_2}$$

$$+ \left( -\frac{y}{z} \right) \frac{\partial y}{\partial t_2} + \left( \frac{z}{x} \right) \frac{\partial y}{\partial t_3} = 0$$

Hence  $\left[ x \frac{\partial y}{\partial x} + y \frac{\partial y}{\partial y} + z \frac{\partial y}{\partial z} = 0 \right]$  proved.

Q. (2) (b)

$$\text{Prove that } \beta(m, n) = \frac{\Gamma(m+n)}{\Gamma(m)\Gamma(n)}, m, n > 0$$

Sol:- we have by property of Gamma function

$$\Gamma(m) = \int_0^\infty e^{-zx} z^{m-1} dx \quad \text{--- (1)}$$

True

$$\Gamma(m) = \int_0^\infty e^{-zx} z^m x^{m-1} dx$$

Multiplying  $e^z z^{m-1}$  both sides and integrate with respect to  $z$  up to limit  $z=0$  to  $z=\infty$

$$\Gamma(m) \int_0^\infty e^{-zx} z^{m-1} dz = \int_0^\infty \int_0^\infty e^{-zx} z^m z^{m-1} dx \cdot e^z z^{m-1} dz$$

$$\Gamma(m) \cdot \Gamma(m) = \int_0^\infty e^{-(1+x)z} z^{m+m-1} dz \cdot \int_0^\infty x^{m-1} dx$$

$$\Gamma(m) \cdot \Gamma(m) = \int_0^\infty e^{-(1+x)z} z^{m+m-1} dz \cdot \int_0^\infty x^{m-1} dx$$

$$m \cdot \overline{m} = \int_{1+x}^{\infty} x^{m-1} dx$$

$$\begin{aligned} m \cdot \overline{m} &= \int_{1+x}^{\infty} x^{m-1} dx \\ &= \int_0^{\infty} x^{m-1} e^{-x} dx \end{aligned}$$

$$m \cdot \overline{m} = \int_0^{\infty} x^{m-1} e^{-x} dx$$

$$(1+x)^{m+n}$$

$$\int_0^{\infty} x^{m+n} = \int_0^{\infty} x^m \cdot x^n = \int_0^{\infty} x^m \cdot \beta(m, n)$$

$$\beta(m, n) = \int_0^{\infty} x^{m+n} dx$$

$$\beta(m, n) = \int_0^{\infty} x^{m+n} dx$$

Ans.

Beispiel

(16)

Q(2) C Find the divergence and curl of the vector

$$\vec{F} = \nabla (x^3 + y^3 + z^3 - 3xyz)$$

Sol:- given  $\vec{F} = \nabla (x^3 + y^3 + z^3 - 3xyz)$

$$= \left( \frac{\partial \hat{i}}{\partial x} + \frac{\partial \hat{j}}{\partial y} + \frac{\partial \hat{k}}{\partial z} \right) (x^3 + y^3 + z^3 - 3xyz)$$

$$\vec{F} = (3x^2 - 3yz)\hat{i} + (3y^2 - 3zx)\hat{j} + (3z^2 - 3xy)\hat{k}$$

$$\text{Then } \operatorname{Div} \vec{F} = \vec{\nabla} \cdot \vec{F} = \left( \frac{\partial \hat{i}}{\partial x} + \frac{\partial \hat{j}}{\partial y} + \frac{\partial \hat{k}}{\partial z} \right) \cdot \vec{F} \quad \hat{i} \cdot \hat{i} = \hat{j} \cdot \hat{j} = \hat{k} \cdot \hat{k} \\ = 1$$

$$= \frac{\partial}{\partial x} (3x^2 - 3yz) + \frac{\partial}{\partial y} (3y^2 - 3zx) + \frac{\partial}{\partial z} (3z^2 - 3xy)$$

$$= 6x + 6y + 6z.$$

$$= 6(x+y+z)$$

Amb.

$$\text{Now } \operatorname{curl} \vec{F} = \vec{\nabla} \times \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 3x^2 - 3xy & 3y^2 - 3xz & 3z^2 - 3xy \end{vmatrix}$$

$$= \hat{i} \left( \frac{\partial}{\partial y} (3z^2 - 3xy) - \frac{\partial}{\partial z} (3y^2 - 3xz) \right) - \hat{j} \left( \frac{\partial}{\partial x} (3z^2 - 3xy) - \frac{\partial}{\partial z} (3x^2 - 3yz) \right) + \hat{k} \left( \frac{\partial}{\partial x} (3y^2 - 3xz) - \frac{\partial}{\partial y} (3x^2 - 3yz) \right)$$

$$\operatorname{curl} \vec{F} = \vec{\nabla} \times \vec{F} = \hat{i} (-3x + 3y) - \hat{j} (-3y + 3z) + \hat{k} (-3z + 3x)$$

$$= 0\hat{i} - 0\hat{j} + 0\hat{k}$$

$$= 0$$

Hence  $\operatorname{curl} \vec{F} = 0$  so that  $\vec{F}$  is irrotational vector field.

Ans.

(P)

$$(d) \quad \text{If } u = f(x) \quad \text{where } x^2 = x^2 + y^2$$

Prove that

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = f''(x) + \frac{1}{x} f'(x)$$

Sol:- given  $u = f(x) \rightarrow x^2 = x^2 + y^2$

True Partial Diff w.r.t x

$$① - \frac{\partial}{\partial x} x \cdot (x) = x \cdot \frac{\partial}{\partial x} x - x \cdot x = f'(x) \cdot x$$

again P. diff. w.r.t y

$$\frac{\partial}{\partial y} x \cdot (x) = x \cdot \frac{\partial}{\partial y} x + x \cdot x = x \cdot 0 + x \cdot x = x^2$$

$$x \cdot (x)'' = x \cdot f''(x) + x \cdot f'(x) \cdot x = x^2 f''(x) + x^2 f'(x)$$

$$x^2 f''(x) + x^2 f'(x) = x^2 \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right)$$

$$x^2 \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right) = f'(x) \int x^2 - x^2 f''(x) =$$

(2)

Ans

formally

$$(x)_{11} f + \frac{x}{T} + (x)_{11} f = \frac{xe}{h\varepsilon} + \frac{xe}{h\varepsilon}$$

$$(x)_{11} f + \left[ \frac{e^x}{x} \right] (x)_{11} f =$$

$$xk \cdot \frac{e^x}{x} + \left[ x - \frac{e^x}{x} \right] (x)_{11} f = \frac{xe}{h\varepsilon} + \frac{xe}{h\varepsilon}$$

$$x = h + x$$

$$\left( h + x \right) (x)_{11} f + \left[ (h + x) - \frac{e^x}{x} \right] (x)_{11} f = \frac{xe}{h\varepsilon} + \frac{xe}{h\varepsilon}$$

(S) and (B) adding

$$2 - (x)_{11} f - \left[ x - \frac{e^x}{x} \right] (x)_{11} f = \frac{xe}{h\varepsilon} - \frac{xe}{h\varepsilon}$$

(B)

Similarly

(e)

Find the extreme value of  $x^2 + y^2 + z^2$  subject to  
 The condition  $ax + by + cz = p$

Sol:- given  $f(x, y, z) = x^2 + y^2 + z^2$   
 and

$$\text{Subject to } \phi = ax + by + cz - p = 0$$

We have by Lagrange's Multiplier Method.  
 we have by Lagrange's eqn.

$$\frac{\partial f}{\partial x} + \lambda \frac{\partial \phi}{\partial x} = 0 \quad \text{---(1)}$$

$$\frac{\partial f}{\partial y} + \lambda \frac{\partial \phi}{\partial y} = 0 \quad \text{---(2)}$$

$$\frac{\partial f}{\partial z} + \lambda \frac{\partial \phi}{\partial z} = 0 \quad \text{---(3)}$$

where  $\lambda$  is the Lagrange's Multiplier.

$$\text{Now } \frac{\partial f}{\partial x} = 2x, \quad \frac{\partial f}{\partial y} = 2y, \quad \frac{\partial f}{\partial z} = 2z.$$

$$\text{and } \frac{\partial \phi}{\partial x} = a, \quad \frac{\partial \phi}{\partial y} = b, \quad \frac{\partial \phi}{\partial z} = c$$

$$\text{Then } 2x + a\lambda = 0 \quad \dots \textcircled{1}$$

$$2y + b\lambda = 0 \quad \dots \textcircled{2}$$

$$2z + c\lambda = 0 \quad \dots \textcircled{3}$$

$$x = \frac{-a\lambda}{2} \quad y = \frac{-b\lambda}{2} \quad z = \frac{-c\lambda}{2}$$

Putting these values in  
 $ax + by + cz = p$

Then

$$\frac{-a^2\lambda}{2} - \frac{b^2\lambda}{2} - \frac{c^2\lambda}{2} = p$$

$$\left[ \begin{array}{l} \lambda = -\frac{2p}{a^2+b^2+c^2} \\ \end{array} \right]$$

Then

$$x = \frac{-a}{2} \times \frac{-2p}{a^2+b^2+c^2} = \frac{ap}{a^2+b^2+c^2}$$

$$y = \frac{bp}{a^2+b^2+c^2}$$

$$z = \frac{cp}{a^2+b^2+c^2}$$

$$\left[ \begin{array}{l} x = \frac{ap}{a^2+b^2+c^2} \\ y = \frac{bp}{a^2+b^2+c^2} \\ z = \frac{cp}{a^2+b^2+c^2} \\ \end{array} \right]$$

These are the extreme values of

$$f(x, y, z) = x^2 + y^2 + z^2 - \frac{a^2p^2}{(a^2+b^2+c^2)^2} + \frac{b^2p^2}{(a^2+b^2+c^2)^2} + \frac{c^2p^2}{(a^2+b^2+c^2)^2} = \frac{(a^2+b^2+c^2)p^2}{(a^2+b^2+c^2)^2} = \frac{p^2}{(a^2+b^2+c^2)}$$

Q. 2) (f)

If  $f(x, y) = \begin{cases} x^3 - y^3 & \text{when } x \neq 0, y \neq 0 \\ 0 & \text{when } x = y = 0 \end{cases}$

Then discuss the continuity of  $f(x, y)$  at the origin.

Sol: — a function  $f(x, y)$  is continuous at origin.

To

$$\lim_{(x,y) \rightarrow (0,0)} f(x, y) = f(0, 0) \quad \dots \quad \textcircled{1}$$

Then,

$$\text{Part - I.} \quad \lim_{\substack{x \rightarrow 0 \\ y \rightarrow 0}} \left( \frac{x^3 - y^3}{x^2 + y^2} \right) = \lim_{\substack{x \rightarrow 0 \\ y \rightarrow 0}} \left\{ \frac{-y^3}{x^2 + y^2} \right\} = 0$$

$$\text{Part - II} \quad \lim_{\substack{x \rightarrow 0 \\ y \rightarrow 0}} \left( \frac{x^3 - y^3}{x^2 + y^2} \right) = \lim_{x \rightarrow 0} \left\{ \frac{x^3}{x^2 + y^2} \right\} = \lim_{x \rightarrow 0} \left\{ x \right\} = 0$$

along the Part  $y = mx$  and  $y = mx^2$ .

$$\lim_{\substack{y \rightarrow mx \\ x \rightarrow 0}} \left( \frac{x^3 - y^3}{x^2 + y^2} \right) = \lim_{x \rightarrow 0} \left\{ \frac{x^3 - m^3 x^3}{x^2 + m^2 x^2} \right\} = \lim_{x \rightarrow 0} \left\{ \frac{x^3 (1 - m^3)}{x^2 (1 + m^2)} \right\} = 0$$

(12)

$$\lim_{\substack{y \rightarrow mx^2 \\ x \rightarrow 0}} \begin{cases} x^3 - y^3 \\ x^2 + y^2 \end{cases} = \lim_{x \rightarrow 0} \begin{cases} x^3 - m^3 x^6 \\ x^2 + m^2 x^4 \end{cases} = \lim_{x \rightarrow 0} \begin{cases} x^3 (1 - m^3 x^3) \\ x^2 (1 + m^2 x^2) \end{cases} = 0$$

Now limit of  $f(x,y)$  along all the paths is same and which is equal to the value of the function at origin. So the function is continuous.

$$\text{i.e. } \lim_{(x,y) \rightarrow (0,0)} f(x,y) = 0 = f(0,0)$$

Ans.

[Part-C]

(B)

Q. ③ ②

FInd The volume and surface area of solid generated by revolving the curve about  $x$ -axis.

$$x = a \cos^3 t, y = a \sin^3 t$$

Sol:- Given eqn of curve. (Astroid)

$$x = a \cos^3 t, y = a \sin^3 t$$

Parameteric form,  $t$  as a parameter.

$$\frac{dx}{dt} = 3a \cos^2 t \cdot (-\sin t)$$

$$(t = \pi/2)$$

$$\frac{dy}{dt} = 3a \sin^2 t \cdot \cos t$$

$$t = \pi$$

$$(-a, 0)$$

$$(0, a)$$

$$(a, 0)$$

$$t = 0$$

curve.

Surface:

$$S = \int_a^b 2\pi y \, ds$$

$$t = -3\pi/2$$

$$= \int_a^b 2\pi y \frac{ds}{dt} \, dt$$

$$\frac{ds}{dt} = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}$$

$$\text{Required surface } S = 2\pi \int_0^{\pi/2} 2\pi y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} \, dt$$

$$S = 4\pi \int_{0}^{\pi/2} a \sin^3 t \cdot \int q^2 \cos^2 t \cdot \sin^2 t + q^2 \sin^2 t \cdot \cos^2 t dt$$

$$S = 4\pi \int_0^{\pi/2} 3a^2 \sin^3 t \cdot \sin^2 t \cdot \int \cos^2 t + \sin^2 t dt$$

$$= 12\pi a^2 \int_0^{\pi/2} \sin^4 t \cdot \cos^2 t dt$$

$$\sin t = u$$

$$\cos t dt = du$$

$$S = 12\pi a^2 \int_0^1 u^4 \cdot du = 12\pi a^2 \left[ \frac{u^5}{5} \right]_0^1$$

$$\boxed{S = \frac{12\pi a^2}{5}}$$

Ans.

Volume

$$V = 2 \times \int_0^{\pi/2} \pi y^2 \frac{dx}{dt} dt = 2 \times \int_0^{\pi/2} \pi (a \sin^3 t)^2 (-3a) \cos^2 t \sin t dt$$

$$V = -6\pi a^3 \int_0^{\pi/2} \sin^6 t \cos^2 t dt = -6\pi a^3 \int_0^{\pi/2} \sin^7 t \cos^2 t dt$$

$$= -6\pi a^3 \times \frac{\sqrt{7+1}}{2} \frac{\sqrt{2+1}}{2} = -3\pi a^3 \times 13 \times \frac{\sqrt{3}}{2} = \frac{21\sqrt{7+2+2}}{2}$$

(20)

$$V = -3\pi a^3 \times (3 \times \frac{\sqrt{3}}{2})$$

$$\Rightarrow -3\pi a^3 \times \cancel{6}^2 \times 16$$

$$\frac{9}{2} \times \frac{7}{2} \times \frac{5}{2} \times \frac{3}{2} \times \frac{\sqrt{3}}{2}$$

3

$$V = -32\pi a^3$$

Ans.

(Neglecting -ve sign)

$$\left\{ V = 32\pi a^3 \right\}$$

105

Q. ③ b)

(ii)

Evaluate

$$I = \int_0^\infty \frac{dx}{1+x^4}$$

$$\text{Let } x^2 = \tan \theta \quad x = \sqrt{\tan \theta}$$

$$2x dx = \sec^2 \theta d\theta$$

$$= \int_0^{\pi/2} \frac{1}{1+\tan^2 \theta} \cdot \frac{\sec^2 \theta}{2\sqrt{\tan \theta}} d\theta$$

$$\frac{1}{2} \int_0^{\pi/2} \frac{1}{\sec^2 \theta} \cdot \frac{\sec^2 \theta}{\sqrt{\tan \theta}} d\theta = \frac{1}{2} \int_0^{\pi/2} \frac{1}{\sqrt{\tan \theta}} d\theta$$

$$= \frac{1}{2} \int_0^{\pi/2} (\tan \theta)^{-1/2} d\theta = \frac{1}{2} \int_0^{\pi/2} (\cot \theta)^{1/2} d\theta = \frac{1}{2} \int_0^{\pi/2} \cos^{-1/2} \theta \sin \theta d\theta$$

$$\rho = \gamma_2 \quad \varphi = -\gamma_2$$

22

we have  $\int_{\theta=0}^{\pi/2} \sin^p \theta \cos^q \theta d\theta$

2

$$\int \frac{P_f}{2} \cdot \sqrt{\frac{Q_f}{2}}$$

## Интегрирование

$$\int_0^{\pi/2} r^2 \cos^2 \theta \sin^2 \theta d\theta \quad P = -r/2 \quad Q = r/2$$

$$= \frac{1}{2} \frac{\int_{-\frac{1}{2}}^{\frac{1}{2}} \int_{-\frac{1}{2}}^{\frac{1}{2}}}{2 \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{1}{2}} = \frac{\int_{-\frac{1}{2}}^{\frac{1}{2}}}{4 \int_{-\frac{1}{2}}^{\frac{1}{2}}} \quad \because r = 1$$

$$\int_{-1}^1 \int_{-1}^1 = \frac{\pi}{4}$$

$$= \frac{1}{4} \left| \frac{1}{4} \right| \left| \frac{3}{4} \right| = \frac{1}{4} \times \frac{\pi}{4} = \frac{1}{16} \pi$$

$$= \frac{1}{4} \left| \frac{1}{4} \right| \left| \frac{1}{4} \right| = \frac{1}{4} \times \frac{1}{4} = \frac{1}{16}$$

$$= \frac{4 \times \frac{1}{16}}{2\sqrt{2}} = \frac{\pi}{2\sqrt{2}} \text{ Ans. or } \frac{\pi\sqrt{2}}{4} \text{ Ans.}$$

Evaluate

(23)

$$(i) I = \int_0^{\pi/2} 16 \cos^4 \theta \sin^2 \theta d\theta \quad \text{Let } 3\theta = t \\ d\theta = \frac{dt}{3}$$

$$= \int_0^{\pi/2} 16 \cos^4 t \cdot \sin^2 t \cdot \frac{dt}{3}$$

$$= \frac{1}{3} \int_0^{\pi/2} 16 \cos^4 t \cdot (2 \sin t \cdot \cos t)^2 dt$$

$$= \frac{4}{3} \int_0^{\pi/2} 16 \cos^6 t \sin^2 t dt$$

$$P=2 \quad q=6$$

$$= \frac{4}{3} \left| \frac{1}{2} \right| \left| \frac{6+1}{2} \right|$$

$$= \frac{4}{3} \times \frac{2}{3} \left| \frac{2+6+2}{2} \right|$$

$$= \frac{2}{3} \times \left| \frac{3}{2} \right| \left| \frac{7}{2} \right|$$

$$= \frac{2}{3} \times \frac{1}{2} \times \frac{5}{2} \times \frac{3}{2} \times \frac{1}{2} \times \frac{1}{4} = \frac{5\pi}{24}$$

Ans.

2

Engineering

Q. (3) C If  $u = f(x, y)$  where  $x = \rho \cos \theta$  and  $y = \rho \sin \theta$

$$y = \rho \sin \theta + \rho \cos \theta$$

Then prove that

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = \frac{\partial^2 u}{\partial \rho^2} + \frac{2u}{\rho^2}$$

Sol:-  $u = f(x, y)$   $x = \rho \cos \theta - \rho \sin \theta$   $\frac{\partial x}{\partial \rho} = \cos \theta$

we have.

$y = \rho \sin \theta + \rho \cos \theta$

$$\frac{\partial y}{\partial \rho} = \sin \theta$$

$\sin \theta = -\sin \theta$

$$\frac{\partial x}{\partial \theta} = \cos \theta$$

$$\frac{\partial y}{\partial \theta} = \sin \theta$$

$$\frac{\partial u}{\partial x} = \frac{\partial u}{\partial \rho} \cdot \frac{\partial \rho}{\partial x} + \frac{\partial u}{\partial \theta} \cdot \frac{\partial \theta}{\partial x}$$

$$\frac{\partial u}{\partial y} = \frac{\partial u}{\partial \rho} \cdot \frac{\partial \rho}{\partial y} + \frac{\partial u}{\partial \theta} \cdot \frac{\partial \theta}{\partial y}$$

$$\frac{\partial u}{\partial x} = \frac{\partial u}{\partial \rho} \cdot \frac{\partial \rho}{\partial x} + \frac{\partial u}{\partial \theta} \cdot \frac{\partial \theta}{\partial x}$$

$$\frac{\partial u}{\partial y} = \frac{\partial u}{\partial \rho} \cdot \frac{\partial \rho}{\partial y} + \frac{\partial u}{\partial \theta} \cdot \frac{\partial \theta}{\partial y}$$

$$\frac{\partial u}{\partial x} = \frac{\partial u}{\partial \rho} \cdot \frac{\partial \rho}{\partial x} + \frac{\partial u}{\partial \theta} \cdot \frac{\partial \theta}{\partial x}$$

$$\frac{\partial u}{\partial y} = \frac{\partial u}{\partial \rho} \cdot \frac{\partial \rho}{\partial y} + \frac{\partial u}{\partial \theta} \cdot \frac{\partial \theta}{\partial y}$$

$$\frac{\partial u}{\partial x} = -\sin \theta + \cos \theta$$

$$\frac{\partial u}{\partial y} = \cos \theta + \sin \theta$$

(25)

$$\frac{\partial^2 u}{\partial y^2} = \frac{\partial}{\partial y} \left( \frac{\partial u}{\partial y} \right) = \left( \frac{\partial}{\partial x} \cdot \cos x + \frac{\partial}{\partial y} \sin x \right) \left( \frac{\partial u}{\partial x} \cos x + \frac{\partial u}{\partial y} \sin x \right)$$

$$= \cos^2 x \frac{\partial^2 u}{\partial x^2} + \sin x \cos x \frac{\partial^2 u}{\partial x \partial y} + \cos x \sin x \frac{\partial^2 u}{\partial y \partial x} + \sin^2 x \frac{\partial^2 u}{\partial y^2}$$

$$= \cos^2 x$$

$$\frac{\partial^2 u}{\partial x^2} + 2 \sin x \cos x \frac{\partial^2 u}{\partial x \partial y} + \sin^2 x \frac{\partial^2 u}{\partial y^2} = \text{---} \quad (3)$$

$$\frac{\partial^2 u}{\partial y^2} = \frac{\partial}{\partial y} \left( \frac{\partial u}{\partial y} \right) = \left( -\sin x \frac{\partial}{\partial x} + \cos x \frac{\partial}{\partial y} \right) \left( -\sin x \frac{\partial u}{\partial x} + \cos x \frac{\partial u}{\partial y} \right)$$

$$= \sin^2 x \frac{\partial^2 u}{\partial x^2} - \sin x \cos x \frac{\partial^2 u}{\partial x \partial y} - \sin x \cos x \frac{\partial^2 u}{\partial y \partial x} + \cos^2 x \frac{\partial^2 u}{\partial y^2}$$

$$= \sin^2 x \frac{\partial^2 u}{\partial x^2} - 2 \sin x \cos x \frac{\partial^2 u}{\partial x \partial y} + \cos^2 x \frac{\partial^2 u}{\partial y^2} = \text{---} \quad (4)$$

Adding eqn (3) and (4)

$$\frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial x^2} = \frac{\partial^2 u}{\partial x^2} \left( \cos^2 x + \sin^2 x \right) + \frac{\partial^2 u}{\partial y^2} \left( \sin^2 x + \cos^2 x \right)$$

$$= \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2}$$

Answer